

# Longitudinal relationship between onset of physical symptoms and functional impairment

Lisa M. McAndrew<sup>1,2</sup>  · Drew A. Helmer<sup>1,3</sup> · Shou-En Lu<sup>1,4</sup> · Helena K. Chandler<sup>1</sup> · Sarah Slotkin<sup>2</sup> · Karen S. Quigley<sup>5,6</sup>

Received: January 10, 2018 / Accepted: May 15, 2018 / Published online: June 9, 2018

© This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2018

**Abstract** Patients with chronic physical symptoms (e.g., chronic pain) often have significant functional impairment (i.e., disability). The fear avoidance model is the dominant theoretical model of how the relationship between chronic physical symptoms and functional impairment develops and proposes a cyclical/bidirectional relationship. There has never been a definitive test of the proposed bi-directional relationship. The current study followed 767 Operation Enduring Freedom/Operation Iraqi Freedom soldiers from pre-deployment, when they were relatively healthy, to 1 year after deployment, when it was anticipated that symptoms would increase or develop. Over the four assessment time points, physical symptom severity consistently predicted worse functional impairment at the

subsequent time point. Functional impairment did not show a consistent relationship with worsening of physical symptom severity. These findings suggest that changes to functional impairment do not have a short-term impact on physical symptom severity.

**Keywords** Disability · Health function · Chronic pain · Medically unexplained symptoms · Iraq · Veteran · Fear avoidance model · Health status · Fatigue

## Introduction

Patients with chronic physical symptoms (e.g., chronic pain) often have significant functional impairment (i.e., disability or limitations in the ability to perform daily and social activities). Cross-sectional studies find that physical symptoms and functional impairment have a strong association. For example, a cross-sectional study of physical symptoms and functional impairment found each accounted for 35% of the variance in the other (Kroenke et al., 1994). This relationship is significant when controlling for mental health variables [e.g., depression, PTSD; (Helmer et al., 2009; Jackson et al., 2006; Tomenson et al., 2013)] and among patients with medically unexplained symptoms/illnesses (Hoffman & Dukes, 2008) and medically explained symptoms/illnesses (Sogutlu et al., 2011) leading some to conclude that chronic physical symptoms and functional impairment are inextricably linked.

The fear avoidance model, a cognitive behavioral model, is the dominant theoretical model of the relationship between chronic physical symptoms and functional impairment and has guided our understanding of this relationship for the past three decades (Lethem et al., 1983). The fear avoidance model proposes a cyclical/

The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs, the Department of Defense or the United States Government.

✉ Lisa M. McAndrew  
lisa.mcandrew@va.gov

<sup>1</sup> War Related Illness and Injury Study Center, Veterans Affairs - New Jersey Health Care System, 385 Tremont Ave, East Orange, NJ 07081, USA

<sup>2</sup> Department of Educational and Counseling Psychology, University at Albany, Albany, USA

<sup>3</sup> New Jersey Medical School, Rutgers University, Newark, USA

<sup>4</sup> School of Public Health, Rutgers University, Piscataway, USA

<sup>5</sup> Department of Veterans Affairs, Bedford Memorial Hospital, Bedford, USA

<sup>6</sup> Interdisciplinary Affective Science Laboratory, Northeastern University, Boston, USA

bidirectional relationship between physical symptoms and functional impairment (Crombez et al., 2012; Wideman et al., 2013). According to the model, acute physical symptoms are hypothesized to trigger pain catastrophizing (a cognitive error in which there is exaggeration of the consequences of an event) which causes fear and avoidance of activity. Avoidance of activity leads to functional impairment and physical deconditioning. Functional impairment (and associated physical deconditioning) then increases the severity of the physical symptoms. The fear avoidance model is also the basis for the dominant behavioral treatment for chronic physical symptoms, cognitive behavioral therapy (Wideman et al., 2013). In cognitive behavioral therapy, patients are encouraged to challenge catastrophizing beliefs and to slowly increase activity to reduce physical symptoms and functional impairment (Brown, 2004; Castell et al., 2011).

Recently, however, there have been questions about the directionality and cyclical nature of the fear avoidance model (Wideman et al., 2013). Studies on the direction of the relationships between variables within the fear avoidance model suggest that common assumptions may be incorrect (Sullivan et al., 2009; Wideman & Sullivan, 2011). For example, two recent prospective studies failed to find that pain catastrophizing precedes fear of activity (Bergbom et al., 2012; Wideman et al., 2009). The cyclical relationship between physical symptoms and functional impairment is a foundational assumption of the fear avoidance model, yet there has not been a study that examines this *bidirectional* relationship.

There are studies that examine one direction of this relationship. These studies find longitudinal evidence that physical symptoms lead to increases in the severity of functional impairment (Momsen et al., 2014; Tomenson et al., 2013). There is conflicting evidence, however, that functional impairment leads to increases in the severity of physical symptoms. Two prospective studies found that functional impairment was associated with greater physical symptoms over time (Bergbom et al., 2012; Bergman et al., 2004) while two reviews found limited evidence that physical deconditioning increases physical symptoms (Smeets et al., 2006; Verbunt et al., 2003). In the fear avoidance model physical deconditioning is proposed to explain the relationship between functional impairment and physical symptoms.

The goal of the current study is to examine the prospective bidirectional (cross-lagged) relationship between physical symptoms and functional impairment. We examined this relationship among Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) military personnel who were first assessed before deployment at a time when we expected to see relatively few physical symptoms and minimal functional impairment, and then

assessed again immediately after deployment, 3 months after deployment and 1 year after deployment. Combat deployment leads to chronic physical symptoms for approximately 30% of Veterans (McAndrew et al., 2016). By examining the cross-lagged relationships between physical symptoms and functional impairment among deploying military personnel, we can better understand these relationships as physical symptoms and functional impairment are developing.

## Methods

### Participants

Participants were recruited as part of the HEROES Project, a prospective longitudinal study of Army National Guard and Army Reserve soldiers deploying to Operation Iraqi/Enduring Freedom. Details on the HEROES project including response bias and drop out can be found in our prior published work using this sample (McAndrew et al., 2016; McAndrew et al., 2013; 2017a, b; Yan et al., 2013). Military personnel between the ages of 18 and 60 were eligible. Exclusion criteria included (a) high blood pressure, (b) medications that produced cardiovascular or autonomic effects, (c) self-reported depression, schizophrenia, or bipolar disorder, or (d) pregnancy. These were specified because they are known to impact physical symptoms and the physiological measures obtained in the study (the latter are not reported here).

### Procedures

The HEROES study was a longitudinal study of military personnel from before to 1 year after deployment. Soldiers were asked to complete questionnaires at four time points: (a) at before deployment while at the Army installation (Time 1; n = 767), (b) immediately upon or within a few days of return from deployment at the Army installation or through the mail if they did not return to the base from which they deployed (Time 2; n = 422), (c) 3-months after deployment through mail (Time 3; n = 286), and (d) 1-year after deployment through mail (Time 4; n = 335). Physical symptom severity at baseline was not related to the likelihood of being lost to follow-up at 1 year post-deployment ( $\chi^2 = 0.80, p = 0.85$ ).

Soldiers were recruited during their deployment readiness medical processing. Study personnel emphasized the voluntary nature of participation and provided study information. At the start of the study 795 enrolled soldiers were eligible; 28 were excluded from analysis because they did not deploy, were officers, or were killed in action. At all-time points, soldiers with significant physical or psy-

chological health concerns were offered referrals to appropriate care providers. Participants were not able to be compensated for their participation while on active duty (Time 1 and 2), but those not on active duty at Time 3 and 4 were reimbursed for their time and effort. All study protocols were approved by multiple Institutional Review Boards and other relevant research committees.

## Measures

### Physical symptoms

Physical symptom severity was measured using the Patient Health Questionnaire-15 (PHQ-15), a widely used self-report questionnaire that measures the presence of common physical health symptoms and their severity (Kroenke et al., 2002). Participants were asked the extent to which they were bothered by each of 15 symptoms listed (e.g., stomach pain; back pain; pain in arms, legs, or joints; menstrual cramps; headaches; chest pain) during the past 7 days. Response options were 0 (*not bothered at all*), 1 (*bothered a little*), and 2 (*bothered a lot*) and responses were added to create a summary score. Participants responded to the PHQ-15 at all four time points.

### Functional impairment

Functional impairment was measured using the mental health composite score (MCS) and physical health composite score (PCS) from the Veterans Rand-36 (VR-36; (Kazis et al., 2004), which was derived from the widely used MOS Short-Form 36 [SF-36 (Ware & Sherbourne, 1992)]. For the purposes of this paper, we will refer to the measure as the SF-36. The mental health functional impairment score (i.e., MCS) captured the degree to which participants experienced problems in social roles and activities because of emotional problems over the past four weeks. The physical health functional impairment score (i.e., PCS) captured the impact of participant's physical health problems on their daily roles and activities. The composite scores were calculated using a standard algorithm. Scores were reported as T-scores ( $M = 50$ ,  $SD = 10$ ), with lower scores indicating more functional impairment. According to the norms, a 2–3 point difference is considered clinically meaningful. Use of the SF-36 as a measure of functional impairment has been demonstrated to have construct validity through comparisons with conceptually similar measures (McHorney et al., 1994).

## Analyses

Descriptive statistics of the sample and Pearson's correlation coefficients for the relationships between physical symptom severity, physical health functional impairment, and mental health functional impairment at all four time points were calculated. We did not impute missing data for the descriptive statistics.

Cross-lagged analyses using a series of linear regression models was used to allow for multiple imputation to handle missing data (Raghunathan et al., 2002; Schafer, 1999). We controlled for age and gender due to previous research indicating that they are significant factors related to symptom reporting (Tomenson et al., 2013). All variables included in the cross-lagged analyses were standardized to a mean of 0 and variance of 1 so that results of the regressions could be reported as standardized coefficient estimates.

For example, the following is a description of the cross-lagged analysis equation in which physical symptoms severity (PHQ-15) is being predicted:  $Y_t$  is the physical symptoms severity at Time t,  $Z_t$  is physical health functional impairment (SF-36 PCS) score at Time t, for  $t = 1, 2, 3, 4$ , and  $\varepsilon_Y$  and  $\varepsilon_Z$  represent random errors. For Time  $t = 2, 3, 4$ , we fitted linear regression models:

$$Y_t = \beta_0 + \beta_{t-1} Y_{t-1} + \gamma_{t-1} Z_{t-1} + \varepsilon_Y + \text{effects of covariates (age, gender)}$$

$$\text{and } Z_t = \alpha_0 + \alpha_{t-1} Z_{t-1} + \lambda_{t-1} Y_{t-1} + \varepsilon_Z + \text{effects of covariates (age, gender)}.$$

These equations provided the associations of present physical symptom severity with previous physical symptom severity ( $\beta_{t-1}$ ) and physical health functional impairment ( $\gamma_{t-1}$ ). They also provided the associations of present physical health functional impairment with previous physical symptom severity ( $\alpha_{t-1}$ ) and physical health functional impairment ( $\lambda_{t-1}$ ). Cross-sectional correlations of physical symptom severity and physical health functional impairment at the same time point were calculated using the Pearson's correlation coefficients ( $r$ ) between the residuals from previously described models of  $Y_t$ 's and  $Z_t$ 's. At Time 1, physical symptom severity and physical health functional impairment correlation was calculated by the Pearson's correlation coefficient between residuals from linear regression models that separately fitted  $Y_1$  and  $Z_1$  as a linear function controlled for age and gender.

We used multiple imputation to handle missing data and created 40 imputed datasets using IVEware (Raghunathan et al., 2002). We then conducted the above described cross-lagged analyses on each imputed dataset, and then combined results using the SAS MIANALYSE procedure

(SASv9.4). Sensitivity analysis was conducted using cross-lagged structural equation modeling in SPSS with the subset of the data with complete data. Results were similar (data not shown). We also conducted sensitivity analysis with the subset of participants with greater physical symptom severity and the subset with greater functional impairment and found that the correlations between physical symptom severity and functional impairment were the same as compared to using the complete data (data not shown).

## Results

### Demographics

Our sample was primarily male (89.7%), Caucasian (77.2%; 9.0% African American, 12.4% Hispanic) and Army National Guard (72.2%; 26.6% Army Reserve or 1.4% Active/Other). On average, participants reported 12.7

(1.6 SD) years of education, or some college education. The average age was 28.0 years (8.38 SD). Means and standard deviations of physical symptom severity and functional impairment at all time points are shown in Table 1.

### Descriptive analyses

The bivariate correlations revealed a strong cross-sectional correlation between physical symptom severity, physical health functional impairment and mental health functional impairment at all time points (Table 2).

### Cross-lagged analyses

The cross-lagged analyses allowed us to examine the cross-sectional and longitudinal relationships between the variables in each model (Fig. 1 is physical health functional impairment and physical symptom severity and Fig. 2 is mental health functional impairment and physical symp-

**Table 1** The mean (SD) of physical symptoms, physical health functional impairment and mental health functional impairment across the deployment spectrum

	Before deployment (Time 1)	Immediately after deployment (Time 2)	3 Months after deployment (Time 3)	1 Year after deployment (Time 4)
Physical symptom severity <sup>a</sup>	5.25 (3.93)	7.94 (4.86)	7.67 (5.11)	7.71 (5.35)
Physical health functional impairment <sup>b</sup>	55.50 (5.21)	53.68 (6.93)	52.84 (7.59)	51.49 (8.75)
Mental health functional impairment <sup>b</sup>	47.98 (9.11)	45.71 (10.71)	44.96 (11.49)	44.90 (12.36)

<sup>a</sup> Higher scores indicate greater physical symptom severity


<sup>b</sup> Higher scores indicate less functional impairment

**Table 2** Bivariate correlations between physical symptom severity, mental and physical health functional impairment across the deployment spectrum


	2	3	4	5	6	7	8	9	10	11	12																						
1. Phy. Sx. before	—	.33*	—	.48*	.49*	—	.22*	—	.32*	.46*	—	.15*	—	.27*	—	.40*	—	.16*	—	.23*													
2. Phy. health func. before		1		—	.15*	—	.19*		.34*	—	.02	—	.23*		.30*		.07	—	.17*	—	.29*	.02											
3. Mental health func. before			—	.15*	1		—	.22*		.12		.42*		—	.24*		—	.00		.41*	—	.14*	—	.06	.34*								
4. Physical Sx immediately after				—	.19*	—	.22*	1		—	.35*		.45*		.60*		—	.38*		—	.41		.58*	—	.26*	—	.40						
5. Phy. health func. immediately after					—	.34*		.01		—	.35*	1		—	.20*		—	.29*		.56*		.06	—	.35*		.50*		.10					
6. Mental health func. immediately after						—	.02		.42*		—	.45*		—	.20		1		—	.40*		—	.02		.56*	—	.28*	—	.02	.55*			
7. Phy. Sx. 3 months after							—	.23*		—	.24*		.60*		—	.29*		1		—	.46*		—	.59*		.73*	—	.44*	—	.51*			
8. Phy. health func. 3 months after								—	.30*		—	.00		—	.38*		.56*		1		—	.06		—	.37*		.59*		.19*				
9. Mental health func. 3 months after									—	.07		.41*		—	.41*		.06		.59*		—	.59*		.06	—	.43*		.09		.66*			
10. Phy. Sx. 1 year after										—	.17*		—	.14*		.58*		—	.35*		.73*		—	.37*		—	.43*	1		—	.46*	—	.57
11. Phy. health func. 1 year after											—	.29*		—	.06		—	.26*		.50*		—	.02		.44*		—	1		—	.00		
12. Mental health func. 1 year after												—	.02		.34*		—	.40*		.10		.55*		—	.51*		.19*		.66*		—	.00	

Phy. Sx., physical symptom severity; Phy. health func., physical health functional impairment; Mental health func., mental health functional impairment; Before, before Deployment; After, after deployment; \*correlations significant at  $p < .01$

**Fig. 1** Cross-lagged relationship between physical symptom severity (PHQ15) and physical health functional impairment (PCS). q:  $p = 0.059$ ; \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.0001$ ; cross-lagged analysis was conducted using series of linear regression models adjusting for age and gender. Multiple imputation (40 datasets) was used to handle missing data (IVEware) with results compiled using Proc MIANALYZE in SAS v9.4.  
<sup>a</sup>Pearson correlation coefficient;  
<sup>b</sup>correlation between residuals of regression analyses



**Fig. 2** Cross-lagged relationship between physical symptom severity (PHQ15) and mental health functional impairment (MCS). \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.0001$ ; cross-lagged analysis was conducted using series of linear regression models adjusting for age and gender. Multiple imputation (40 datasets) was used to handle missing data (IVEware) with results compiled using Proc MIANALYZE in SAS v9.4.  
<sup>a</sup>Pearson correlation coefficient;  
<sup>b</sup>correlation between residuals of regression analyses



tom severity). We found a moderate cross-sectional relationship between physical symptom severity and physical health functional impairment at all time points (before deployment:  $r = -.35$ , immediately after deployment  $r = -.28$ , 3 months after deployment  $r = -.32$ , and 1 year after deployment  $r = -.26$ ; Fig. 1). There was a strong cross-sectional relationship between physical symptom severity and mental health functional impairment (before deployment:  $r = -.53$ , immediately after deployment  $r = -.40$ , 3 months after deployment  $r = -.52$ , and 1 year after deployment  $r = -.47$ ; Fig. 2).

There was a strong longitudinal relationship between physical symptom severity at each time point to the subsequent time point. This was true for the model with physical symptom severity and physical health functional impairment (before deployment to immediately after deployment  $\beta = .44$ , immediately after deployment to 3 months after deployment  $\beta = .50$ , 3 months after deployment to 1 year after deployment  $\beta = .63$ ; Fig. 1) and the model with physical symptom severity and mental

health functional impairment (before deployment to immediately after deployment  $\beta = .47$ , immediately after deployment to 3 months after deployment  $\beta = .47$ , 3 months after deployment to 1 year after deployment  $\beta = .63$ ; Fig. 2).

The data revealed a moderate to strong longitudinal relationship between physical health functional impairment at one time point and at the subsequent time point (from before deployment to immediately after deployment  $\lambda = .28$ , from immediately after deployment to 3 months after deployment  $\lambda = .47$  and from 3 months after deployment to 1 year after deployment  $\lambda = .44$ ; see Fig. 1). The data revealed a significant longitudinal relationship between mental health functional impairment at one time point and at the subsequent time point (from before deployment to immediately after deployment  $\lambda = .31$ , from immediately after deployment to 3 months after deployment  $\lambda = .53$  and from 3 months after deployment to 1 year after deployment  $\lambda = .56$ ; see Fig. 2).

Physical symptom severity at each time point showed a consistent small association with worse physical health functional impairment at the subsequent time point (controlling for prior physical symptom severity and prior physical health functional impairment). Physical symptom severity captured before deployment was related to physical health functional impairment ( $\lambda_1 = -.12$ ) and mental health functional impairment ( $\lambda_1 = -.15$ ) after deployment, physical symptom severity immediately after deployment predicted physical health functional impairment ( $\lambda_1 = -.14$ ) and mental health functional impairment ( $\lambda_1 = -.14$ ) 3 months after deployment and physical symptoms 3 months after deployment predicted physical health functional impairment ( $\lambda_1 = -.21$ ) and mental health functional impairment ( $\lambda_1 = -.17$ ) (Figs. 1 and 2).

The cross-lagged result did not show a consistent relationship between either physical or mental health functional impairment to later physical symptom severity (after controlling for prior physical symptom severity or prior functional impairment). Neither physical nor mental health functional impairment before deployment was related to physical symptom severity immediately after deployment. Physical health functional impairment ( $\hat{\gamma}_2 = -.13$ ,  $p = .059$ ) and mental health functional impairment ( $\hat{\gamma}_2 = -.17$ ) immediately after deployment showed a small relationship with physical symptom severity 3 months after deployment (Figs. 1 and 2). Neither physical nor mental health functional impairment 3 months after deployment was related to physical symptom severity 1 year after deployment.

## Discussion

The goal of this study was to examine the bidirectional relationship between physical symptoms and functional impairment proposed by the fear avoidance model. We examined this relationship in a sample of OEF/OIF Veterans who were assessed from before to 1 year after deployment. This allowed us to examine the relationships among a relatively healthy sample as they were developing new or worsening chronic symptoms.

As expected, greater physical symptom severity was prospectively associated with worsening mental and physical health functional impairment. This is consistent with the extant literature which has shown a relationship between having chronic physical symptoms and physical health functional impairment (Kroenke et al., 1994; Tomenson et al., 2013). The current study adds to the literature by examining this relationship using a prospective longitudinal design and by assessing this relationship as physical symptoms are developing. Previous studies mostly

examine this relationship among patients with established chronic symptoms.

There was not a consistent relationship between greater physical or mental health functional impairment and increasing physical symptom severity. This is consistent with a review that found no relationship between physical deconditioning and increases in physical symptoms (Verbunt et al., 2003). Our data suggest that in the short-term, functional impairment likely does not significantly contribute to acute physical symptoms becoming chronic, as is proposed by the fear avoidance model (Leeuw et al., 2007; Wideman et al., 2013).

This study examined the prospective relationship of functional impairment to subsequent physical symptoms over the course of 2 years. The findings do not preclude the possibility that functional impairment may cause increases in physical symptoms over a longer time period. Functional impairment is a predictor of mortality and major morbidity (Afilalo et al., 2012; Idler & Kasl, 1991) and is thought to contribute to obesity, diabetes and heart disease (Bortz, 1984; Kaplan et al., 1996). Obesity, diabetes, and heart disease are all linked to increases in pain and physical symptoms over the course of a lifetime (Calcutt, 2002; Fernandez et al., 2016; Shiri et al., 2009).

Our sample consisted of a relatively healthy population of soldiers and most had average or better health function 1 year after deployment; soldiers with certain, more serious medical conditions were excluded. It is possible that there is a relationship between functional impairment and subsequent physical symptoms among individuals with more severe functional impairment. We explored this possibility, and did not find different results when examining a subset of those with the worst functional impairment or a subset of those with the greatest physical symptom severity as compared to the full sample (data available upon request). This should be explored further in future studies among populations with more medical conditions. Another consideration is our use of a self-report measure of functional impairment. It will be important to examine these relationships using other measures, particularly objective measures, such as the 6 min walk.

The use of a military sample was a strength and a limitation. Deployment is a known predictor of increases in functional impairment and physical symptoms. Thus, studying military personnel allowed us to examine the relationship between these variables while they are developing. Further, Veterans are disproportionately burdened by functional impairment and chronic symptoms, therefore understanding the relationship between these variables in a military sample is particularly important. A limitation is that we do not know how generalizable these findings are. Further, the difficulty following military personnel as they returned from deployment contributed to our high attrition.

The findings of the current study suggest that there is not a bi-directional relationship between physical symptoms and functional impairment while they are developing, rather physical symptoms are associated with decrements in functional impairment and not vice versa. These findings suggest the need to rethink the cyclical nature of the fear avoidance model and to better explore the mechanisms through which acute symptoms become chronic. Cognitive behavioral treatments for disorders characterized by physical symptoms (chronic pain, fibromyalgia) have smaller effect sizes as compared to treatments for other disorders, such as anxiety disorders (Hofmann et al., 2012). A better understanding of the pathways by which acute symptoms become chronic and relate to functional impairment may lead to better treatments and improvements in patient outcomes.

**Acknowledgements** This study was supported by grants from the Department of Veterans Affairs, Health Services Research and Development Service (IIR 0202–296 to K. Quigley; IIRK2HX001369 to L. M. McAndrew); the NJ War Related Illness and Injury Study Center, the NJ REAP (REA 03–021); and the Deployment Health Clinical Center, Walter Reed Army Medical Center.

#### Compliance with ethical standards

**Conflict of interest** Lisa M. McAndrew, Drew A. Helmer, Shou-En Lu, Helena K. Chandler, Sarah Slotkin, and Karen S. Quigley declare that they have no conflict of interest.

**Human and animal rights and Informed consent** All procedures were in accordance with the ethical standards of the institutional research committees and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study. Provided Funding information has to be tagged.

#### References

Afilalo, J., Mottillo, S., Eisenberg, M. J., Alexander, K. P., Noiseux, N., Perrault, L. P., et al. (2012). Addition of frailty and disability to cardiac surgery risk scores identifies elderly patients at high risk of mortality or major morbidity. *Circulation Cardiovascular Quality and Outcomes*, 5, 222–228.

Bergbom, S., Boersma, K., & Linton, S. J. (2012). Both early and late changes in psychological variables relate to treatment outcome for musculoskeletal pain patients at risk for disability. *Behaviour Research and Therapy*, 50, 726–734.

Bergman, S., Jacobsson, L. T., Herrström, P., & Petersson, I. F. (2004). Health status as measured by SF-36 reflects changes and predicts outcome in chronic musculoskeletal pain: A 3-year follow up study in the general population. *Pain*, 108, 115–123.

Bortz, W. M., II. (1984). The disuse syndrome. *Western Journal of Medicine*, 141, 691–694.

Brown, R. J. (2004). Psychological mechanisms of medically unexplained symptoms: An integrative conceptual model. *Psychological Bulletin*, 130, 793–812.

Calcutt, N. A. (2002). Potential mechanisms of neuropathic pain in diabetes. *International Review of Neurobiology*, 50, 205–228.

Castell, B. D., Kazantzis, N., & Moss-Morris, R. E. (2011). Cognitive behavioral therapy and graded exercise for chronic fatigue syndrome: A meta-analysis. *Clinical Psychology: Science and Practice*, 18, 311–324.

Crombez, G., Eccleston, C., Van Damme, S., Vlaeyen, J. W., & Karoly, P. (2012). Fear-avoidance model of chronic pain: The next generation. *The Clinical Journal of Pain*, 28, 475–483.

Fernandez, M., Ordoñana, J. R., Hartvigsen, J., Ferreira, M. L., Refshauge, K. M., Sánchez-Romera, J. F., et al. (2016). Is chronic low back pain associated with the prevalence of coronary heart disease when genetic susceptibility is considered? A co-twin control study of Spanish twins. *PLoS ONE*, 11, e0155194.

Helmer, D. A., Chandler, H. K., Quigley, K. S., Blatt, M., Teichman, R., & Lange, G. (2009). Chronic widespread pain, mental health, and physical role function in OEF/OIF veterans. *Pain Medicine*, 10, 1174–1182.

Hoffman, D., & Dukes, E. (2008). The health status burden of people with fibromyalgia: A review of studies that assessed health status with the SF-36 or the SF-12. *International Journal of Clinical Practice*, 62, 115–126.

Hofmann, S. G., Asnaani, A., Vonk, I. J., Sawyer, A. T., & Fang, A. (2012). The efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognit Ther Res*, 36, 427–440. <https://doi.org/10.1007/s10608-012-9476-1>

Idler, E. L., & Kasl, S. (1991). Health perceptions and survival: Do global evaluations of health status really predict mortality? *Journal of gerontology*, 46, S55–S65.

Jackson, J., Fiddler, M., Kapur, N., Wells, A., Tomenson, B., & Creed, F. (2006). Number of bodily symptoms predicts outcome more accurately than health anxiety in patients attending neurology, cardiology, and gastroenterology clinics. *Journal of Psychosomatic Research*, 60, 357–363.

Kaplan, G. A., Goldberg, D. E., Everson, S. A., Cohen, R. D., Salonen, R., Tuomilehto, J., et al. (1996). Perceived health status and morbidity and mortality: Evidence from the Kuopio ischaemic heart disease risk factor study. *International Journal of Epidemiology*, 25, 259–265.

Kazis, L. E., Lee, A., Spiro, A., III, Rogers, W., Ren, X. S., Miller, D. R., et al. (2004). Measurement comparisons of the medical outcomes study and veterans SF-36 health survey. *Health Care Financing Review*, 25, 43–58.

Kroenke, K., Spitzer, R. L., & Williams, J. B. (2002). The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosomatic Medicine*. <https://doi.org/10.1097/0006842-200203000-00008>

Kroenke, K., Spitzer, R. L., Williams, J. B., Linzer, M., Hahn, S. R., & de Gruy, F. V. (1994). Physical symptoms in primary care. Predictors of psychiatric disorders and functional impairment. *Archives of Family Medicine*. <https://doi.org/10.1001/archfam.3.9.774>

Leeuw, M., Goossens, M. E., Linton, S. J., Crombez, G., Boersma, K., & Vlaeyen, J. W. (2007). The fear-avoidance model of musculoskeletal pain: Current state of scientific evidence. *Journal of Behavioral Medicine*, 30, 77–94.

Lethem, J., Slade, P., Troup, J., & Bentley, G. (1983). Outline of a fear-avoidance model of exaggerated pain perception—I. *Behaviour Research and Therapy*, 21, 401–408.

McAndrew, L. M., D'Andrea, E., Lu, S. E., Abbi, B., Yan, G. W., Engel, C., et al. (2013). What pre-deployment and early post-deployment factors predict health function after combat deployment? A prospective longitudinal study of Operation Enduring Freedom (OEF)/Operation Iraqi Freedom (OIF) soldiers. *Health Quality of Life Outcomes*, 11, 73. <https://doi.org/10.1186/1477-7525-11-73>

McAndrew, L., Helmer, D., Phillips, L., Chandler, H., Ray, K., & Quigley, K. (2016). Iraq and Afghanistan Veterans report symptoms consistent with chronic multisymptom illness one year after deployment. *Journal of Rehabilitation Research and Development*, 53, 59–70. <https://doi.org/10.1682/JRRD.2014.10.0255>

McAndrew, L. M., Markowitz, S., Lu, S.-E., Borders, A., Rothman, D., & Quigley, K. S. (2017a). Resilience during war: Better unit cohesion and reductions in avoidant coping are associated with better mental health function after combat deployment. *Psychological Trauma: Theory, Research, Practice, and Policy*, 9, 52–61. <https://doi.org/10.1037/tra0000152>

McAndrew, L. M., Phillips, L. A., Helmer, D. A., Maestro, K., Engel, C. C., Greenberg, L. M., et al. (2017b). High healthcare utilization near the onset of medically unexplained symptoms. *Journal of Psychosomatic Research*, 98, 98–105. <https://doi.org/10.1016/j.jpsychores.2017.05.001>

McHorney, C. A., Ware, Jr., J. E., Lu, J. R., & Sherbourne, C. D. (1994). The MOS 36-item Short-Form Health Survey (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse patient groups. *Medical Care*, 32, 40–66.

Momsen, A.-M. H., Jensen, O. K., Nielsen, C. V., & Jensen, C. (2014). Multiple somatic symptoms in employees participating in a randomized controlled trial associated with sickness absence because of nonspecific low back pain. *The Spine Journal*, 14, 2868–2876.

Raghunathan, T. E., Solenberger, P. W., & Van Hoewyk, J. (2002). *IVEware: Imputation and variance estimation software*. Ann Arbor, MI: Survey Methodology Program, Survey Research Center, Institute for Social Research, University of Michigan.

Schafer, J. L. (1999). Multiple imputation: A primer. *Statistical Methods in Medical Research*, 8, 3–15.

Shiri, R., Karppinen, J., Leino-Arjas, P., Solovieva, S., & Viikari-Juntura, E. (2009). The association between obesity and low back pain: A meta-analysis. *American Journal of Epidemiology*, 171, 135–154.

Smeets, R. J., Wade, D., Hidding, A., Van Leeuwen, P. J., Vlaeyen, J. W., & Knottnerus, J. A. (2006). The association of physical deconditioning and chronic low back pain: A hypothesis-oriented systematic review. *Disability and Rehabilitation*, 28, 673–693.

Sogutlu, A., Levenson, J. L., McCloud, D. K., Rosef, S. D., & Smith, W. R. (2011). Somatic symptom burden in adults with sickle cell disease predicts pain, depression, anxiety, health care utilization, and quality of life: The PiSCES project. *Psychosomatics*, 52, 272–279.

Sullivan, M., Tanzer, M., Stanish, W., Fallaha, M., Keefe, F. J., Simmonds, M., et al. (2009). Psychological determinants of problematic outcomes following total knee arthroplasty. *Pain*, 143, 123–129.

Tomenson, B., Essau, C., Jacobi, F., Ladwig, K. H., Leiknes, K. A., Lieb, R., et al. (2013). Total somatic symptom score as a predictor of health outcome in somatic symptom disorders. *The British Journal of Psychiatry*. <https://doi.org/10.1192/bjp.bp.112.114405>

Verbunt, J. A., Seelen, H. A., Vlaeyen, J. W., Heijden, G. J., Heuts, P. H., Pons, K., et al. (2003). Disuse and deconditioning in chronic low back pain: Concepts and hypotheses on contributing mechanisms. *European Journal of Pain*, 7, 9–21.

Ware, J. E. S., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36). *Medical Care*, 30, 473–483.

Wideman, T. H., Adams, H., & Sullivan, M. J. (2009). A prospective sequential analysis of the fear-avoidance model of pain. *Pain*, 145, 45–51.

Wideman, T. H., Asmundson, G. G., Smeets, R. J. M., Zautra, A. J., Simmonds, M. J., Sullivan, M. J., et al. (2013). Re-thinking the fear avoidance model: Toward a multi-dimensional framework of pain-related disability. *Pain*, 154, 2262.

Wideman, T. H., & Sullivan, M. J. (2011). Differential predictors of the long-term levels of pain intensity, work disability, healthcare use, and medication use in a sample of workers' compensation claimants. *Pain*, 152, 376–383.

Yan, G. W., McAndrew, L. M., D'Andrea, E. A., Lange, G., Santos, S. L., Engel, C. C., et al. (2013). Self-reported stressors of National Guard women veterans before and after deployment: The relevance of interpersonal relationships. *Journal of General Internal Medicine*, 28, S549–S555. <https://doi.org/10.1007/s11606-012-2247-6>