

BRIEF REPORT

Rumination Moderates the Associations Between PTSD and Depressive Symptoms and Risky Behaviors in U. S. Veterans

Ashley Borders,^{1,2} Lisa M. McAndrew,^{1,3} Karen S. Quigley,^{4,5} and Helena K. Chandler¹

¹Department of Veterans Affairs, NJ War Related Illness & Injury Study Center, East Orange, New Jersey, USA

²The College of New Jersey, Ewing, New Jersey, USA

³New Jersey Medical School-University of Medicine & Dentistry of NJ, Newark, New Jersey, USA

⁴Department of Veterans Affairs, Bedford Memorial Hospital, Bedford, Massachusetts, USA

⁵Interdisciplinary Affective Science Laboratory, Northeastern University, Boston, Massachusetts, USA

Risky behaviors, including unsafe sex, aggression, rule breaking, self-injury, and dangerous substance use have become a growing issue for U.S. veterans returning from combat deployments. Evidence in nonveteran samples suggests that risky behaviors reflect efforts to cope with and alleviate depressive and/or anxious symptoms, particularly for individuals with poor emotion-regulation skills. These associations have not been studied in veterans. Rumination, or repeated thoughts about negative feelings and past events, is a coping strategy that is associated with several psychopathologies common in veterans. In this cross-sectional study, 91 recently returned veterans completed measures of trait rumination, self-reported risky behaviors, and symptoms of posttraumatic stress disorder (PTSD) and depression. Analyses revealed that veterans with more depressive and PTSD symptoms reported more risky behaviors. Moreover, rumination significantly interacted with PTSD symptoms and depressive symptoms (both $\beta = .21$, $p < .05$), such that psychiatric symptoms were associated with risky behaviors only for veterans with moderate to high levels of rumination. Although cross-sectional, these findings support theory that individuals with poor coping skills may be particularly likely to respond to negative mood states by engaging in risky behaviors. Implications include using rumination-focused interventions with veterans in order to prevent engagement in risky behaviors.

Risky behaviors have received recent media attention as a growing issue for U. S. war veterans. Research on individuals following combat deployments to Iraq or Afghanistan found that 47% lose their temper and 15% destroy things in anger (Killgore, Stetz, Castro, & Hoge, 2008). Moreover, they engage in risky recreation (25%), self-harm (10%), and illegal drug use (7%) at higher rates than those who have not yet deployed (Thomsen, Stander, McWhorter, Rabenhorst, & Milner, 2011). Furthermore, greater exposure to combat predicts more risk-taking (Killgore et al., 2008). Risky behaviors lead to incarceration, social isolation, and danger to other individuals.

Therefore, understanding variables that make veterans more likely to engage in risky behaviors is crucial.

Risky behaviors may arise in part from a desire to avoid or escape aversive emotional states (Cooper, Agocha, & Sheldon, 2000). Thus, individuals may binge on substances, steal, or aggress because the immediate reduction of emotional distress is worth any potential future consequence. In support of this theory, significant associations exist between depressive and anxious symptoms and later risky behaviors (Auerbach, Abela, & Ho, 2007; Auerbach, Claro, Abela, Zhu, & Yao, 2010).

Moreover, risky behaviors may be particularly likely for individuals with poor emotion regulation. These individuals may turn to risky behaviors because they lack more effective coping strategies. Emotion regulation motives, in fact, predict risky alcohol and sex behaviors (Cooper et al., 2000). Auerbach and colleagues (Auerbach, Abela, Zhu, & Yao, 2007; Auerbach et al., 2010) also found correlations between poor emotion regulation skills and greater risky behaviors. Moreover, maladaptive emotion regulation interacted with depressive and anxious symptoms to predict increases in risky behaviors. These associations have never been studied in veteran samples, for whom risky behaviors may be particularly likely in response to depression and PTSD. Moreover, identifying specific

This material is the result of work supported with resources and the use of facilities at the War Related Illness and Injury Study Center (WRIISC), Department of Veterans Affairs, New Jersey Health Care System. The views expressed are those of the authors and do not represent the views of the Department of Veterans Affairs or the United States Government.

Correspondence concerning this article should be addressed to Ashley Borders, Department of Veterans Affairs, NJ–War Related Illness and Injury Study Center, East Orange, New Jersey. E-mail: borders@tcnj.edu

Published 2012. This article is a US Government work and is in the public domain in the USA. View this article online at wileyonlinelibrary.com
DOI: 10.1002/jts.21733

maladaptive coping strategies might suggest particular targets for clinical interventions.

We propose that rumination, or thinking repeatedly and passively about negative experiences, is a coping strategy that may put depressed or anxious veterans at particular risk for increased risky behaviors. Rumination is linked to the maintenance of both depression and PTSD (Bennett & Wells, 2010; Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008). Because rumination involves dwelling on negative experiences, it prolongs and intensifies existing negative emotions. Moreover, it is associated with risky behaviors such as aggression, self-harm, and binge drinking in nonveteran samples (Collins & Bell, 1997; Nolen-Hoeksema et al., 2008). Previous studies found associations between rumination and increased emotional distress in Vietnam and World War II veterans (Holman & Silver, 1998; Kraaij & Garnefski, 2006). No study of which we are aware, however, has examined whether rumination is associated with risky behaviors among veterans.

In this cross-sectional study, we examined psychiatric symptoms and rumination as factors that may contribute to risky behaviors in a sample of combat veterans seeking treatment from the Department of Veterans Affairs (VA). We hypothesized that greater PTSD and depressive symptoms would be associated with increased risky behaviors and that these associations would be moderated by rumination.

Method

Participants and Procedure

Veterans were recruited from the VA New Jersey War Related Illness and Injury Study Center (NJ WRIISC), a tertiary care specialty clinic that provides care for veterans with deployment-related health concerns. More than 1,200 veterans participated in comprehensive clinical evaluations at the NJ WRIISC between 2004 and 2010, during which they completed a baseline clinical evaluation. All veterans evaluated at the WRIISC were invited to participate in this follow-up study. Of these, 324 consented and were mailed a follow-up questionnaire packet up to three times (and received two follow-up phone calls); 28% completed the packet. We found no differences in initial PTSD symptoms, age, education, or marital status between those who did and did not complete this survey. Thus, although our response rate was low, we suggest that our current sample was relatively representative of the group from which we sampled. All research was approved by VA New Jersey Healthcare System IRB and appropriate research committees.

Our sample of 91 was 87% male. Their mean age was 37 years ($SD = 11.3$, range = 21–66). Over 90% identified as Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) veterans; the remainder served in a previous conflict. Approximately 39% identified as White, 30% Latino/Latina, 17% Black, 2.5% Asian, 2.5% American Indian, and 9% Other. The average years of education was 14 ($SD = 2.0$, range = 11–18).

Measures

Risky behaviors. The 20-item Risky Behavior Questionnaire for Adolescents (Auerbach & Gardiner, 2012) assesses frequency of engagement in unsafe sex, substance use, and aggressive, dangerous, and illegal behaviors in the past month. Response options ranged from *never* to *4 times or more per week*. This measure correlates strongly with impulsivity and alcohol use, suggesting validity (Auerbach et al., 2007). To make the wording appropriate for adults, we changed “boyfriend or girlfriend” to “significant other” and “skipped class” to “skipped work (or school).” We also omitted three items requiring psychiatric follow-up (e.g., cutting yourself). Higher sum scores indicated more risky behavior (Cronbach’s α in this sample = .86). Due to skipped items, four participants lacked complete data for this measure.

Rumination. The 12-item Rumination subscale of the Rumination and Reflection Questionnaire (Trapnell & Campbell, 1999) assesses the tendency to engage in self-focused repetitive thinking about threats or losses (e.g., “Long after an argument is over with, my thoughts keep going back to what had happened”). Participants indicate level of agreement on a 5-point scale from *strongly disagree* to *strongly agree*. This measure correlates highly with measures of brooding, depression rumination, and thinking repeatedly about negative things (Siegle, Moore, & Thase, 2004). Higher sum scores indicated more rumination ($\alpha = .94$). Three participants lacked complete data.

PTSD symptoms. The 17-item PTSD Checklist (see Blanchard, Jones-Alexander, Buckley, & Forneris, 1996) assesses the presence and frequency of PTSD symptoms over the past month. Participants respond on a 5-point scale from *not at all* to *extremely*. This measure correlates highly with the Clinician Administered PTSD scale (Blanchard et al., 1996). Higher sum scores indicated more PTSD symptomatology ($\alpha = .97$). Two participants lacked complete data.

Depressive symptoms. The Patient Health Questionnaire (PHQ-8; Kroenke et al., 2009) assesses the presence and frequency of eight depressive symptoms over the past 2 weeks, omitting suicidal ideation for research purposes. We included the additional item that assesses degree of functional impairment. Participants respond on a 4-point scale from *not at all* to *nearly every day*. Substantial research supports the construct validity of this measure as a screening and diagnostic tool (Kroenke et al., 2009). Higher sum scores indicated greater depressive symptoms ($\alpha = .92$). Ten participants lacked complete data.

Data Analysis

To investigate the extent that rumination, psychiatric symptoms, and their interaction were related to risky behaviors, we conducted two regression analyses: one with depressive symptoms as the independent variable and one with PTSD symptoms.

Table 1
Descriptive Statistics and Correlations

Variable	Mean	SD	Range	1.	2.	3.
1. Depressive symptoms	12.43	7.70	0–26	—		
2. PTSD symptoms	50.13	20.18	17–85	.85	—	
3. Rumination	43.33	10.94	12–60	.61	.73	—
4. Risky behaviors	7.65	8.76	0–34	.56	.65	.49

Note. $N = 91$. $p < .001$ for all correlations. PTSD = Posttraumatic stress disorder.

Using the SPSS macro PROCESS (Hayes, 2012), we entered centered main effects of symptoms and rumination and their interaction as a single block. We probed significant interactions by estimating the conditional effects of symptoms on risky behaviors at the 10th, 25th, 50th, 75th, and 90th percentiles of rumination (Hayes, 2012). Analyses only included participants with complete data on relevant variables.

Results

Greater rumination significantly correlated with more risky behaviors, PTSD symptoms, and depression symptoms (Table 1). In addition, higher risky behavior correlated with more depressive and PTSD symptoms. Approximately 53% of the sample scored above the cutoff of 50 for PTSD, and 38% reported moderately severe or severe depressive symptoms. In the past month, 20% reported being in a physical fight, 28% bullied a peer, 20% had unsafe sex, 26% drove recklessly, 20% destroyed property, 33% verbally harassed someone, and 22% engaged in acts of revenge.

In regression equations, both depressive and PTSD symptoms emerged as significantly associated with risky behaviors (Table 2). The main effect of rumination was significant for depressive symptoms only. The interactions of rumination with depressive symptoms and rumination with PTSD symptoms, however, were both significant. Analysis of conditional effects

revealed that PTSD symptoms were associated with risky behaviors at the 25th, 50th, 75th, and 90th percentiles but not at the very low level (10th percentile) of rumination. Depressive symptoms was associated with risky behaviors only at moderate, high, and very high levels of rumination (50th, 75th, and 90th percentiles).

Discussion

In line with previous research, veterans with greater depressive and PTSD symptoms reported more risky behaviors. Moreover, these associations appeared to be moderated by rumination. Specifically, PTSD symptoms were associated with risky behaviors for all but very low levels of rumination. Depressive symptoms were only associated with risky behaviors for veterans with moderate to high levels of rumination. Although cross-sectional, these findings are consistent with theory that individuals with poor coping skills may be especially likely to respond to negative moods by engaging in risky behaviors. Future research with veterans should examine these variables longitudinally to establish directional associations.

Very few studies have explicitly measured rumination in veterans. Greater rumination was associated with worse psychiatric symptoms and more risky behaviors. Given the high prevalence of depression and PTSD in combat veterans, rumination

Table 2
Multiple Regression Analyses for Risky Behaviors Using Depressive and PTSD Symptoms Separately

Variable	<i>b</i>	<i>SE b</i>	<i>B</i>	<i>t</i>
Depressive symptoms				
Depressive symptoms	.42	.13	.38	3.14**
Rumination	.26	.10	.31	2.49*
Depression × Rumination	.02	.01	.21	2.01*
Overall model: adjusted $R^2 = .37$, $F(3, 70) = 15.50^{***}$				
PTSD symptoms				
PTSD symptoms	.26	.05	.59	4.89***
Rumination	.13	.11	.17	1.21
PTSD × Rumination	.01	.003	.21	2.21*
Overall model: adjusted $R^2 = .45$, $F(3, 78) = 23.16^{***}$				

Note. N for depressive symptoms analyses = 74. N for PTSD symptoms analyses = 82. PTSD = Posttraumatic stress disorder.

* $p < .05$. ** $p < .01$. *** $p < .001$.

may be an important risk factor to target in this population. Ruminators show poor executive functioning (Whitmer & Banich, 2007), which is also seen in veterans with traumatic brain injuries (Campbell et al., 2009). Therefore, rumination may exacerbate vulnerabilities that veterans are already experiencing, further increasing the likelihood of risky behaviors. Future studies with veteran samples could also examine whether specific types of rumination (e.g., depressive, anger) relate to particular outcomes.

One important limitation was that we used a measure of risky behaviors created for adolescents, and we omitted three items. Therefore, the measure may have had different psychometric properties from previously reported studies. Given the prevalence of risky behaviors in veterans, future research should validate a measure for use with veterans. Additionally, we sampled veterans who sought treatment at a tertiary care VA clinic and who had a low response rate to this survey and some incomplete data. Therefore, our results may not be generalizable to the larger veteran or active duty population and should be replicated.

The finding that psychiatric symptoms were not related to risky behaviors at low levels of rumination suggests that interventions targeting rumination may help prevent dangerous behaviors. Rumination-focused cognitive-behavioral therapy teaches clients alternative coping strategies (e.g., disclosure of emotions, concrete thinking) and effectively decreases symptoms of depression and PTSD (Sezibera, Van Broeck, & Philippot, 2009). Alternatively, training in mindfulness, or intentional and compassionate attention to the present moment, decreases both rumination and delinquent behavior (Shapiro, Oman, Thoresen, Plante, & Flinders, 2008). Given these promising treatments, mental health providers who work with veterans might consider using rumination-focused interventions to help clients cope with negative affective states and potentially prevent risky behaviors.

References

Auerbach, R. P., & Gardiner, C. K. (2012). Moving beyond the trait conceptualization of self-esteem: The prospective effect of impulsiveness, coping, and risky behavior engagement. *Behaviour Research and Therapy*, 50, 596–603. Retrieved from PubMed.

Auerbach, R. P., Abela, J. Z., & Ho, M. (2007). Responding to symptoms of depression and anxiety: Emotion regulation, neuroticism, and engagement in risky behaviors. *Behaviour Research and Therapy*, 45, 2182–2191. doi:10.1016/j.brat.2006.11.002

Auerbach, R. P., Abela, J. Z., Zhu, X., & Yao, S. (2007). A diathesis-stress model of engagement in risky behaviors in Chinese adolescents. *Behaviour Research and Therapy*, 45, 2850–2860. doi:10.1016/j.brat.2007.05.005

Auerbach, R. P., Claro, A., Abela, J. Z., Zhu, X., & Yao, S. (2010). Understanding risky behavior engagement amongst Chinese adolescents. *Cognitive Therapy and Research*, 34, 159–167. doi:10.1007/s10608-009-9238-x

Bennett, H., & Wells, A. (2010). Metacognition, memory disorganization and rumination in posttraumatic stress symptoms. *Journal of Anxiety Disorders*, 24, 318–325. doi:10.1016/j.janxdis.2010.01.004

Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy*, 34, 669–673. doi:10.1016/0005-7967(96)00033-2

Campbell, T. A., Nelson, L. A., Lumpkin, R., Yoash-Gantz, R. E., Pickett, T. C., & McCormick, C. L. (2009). Neuropsychological measures of processing speed and executive functioning in combat veterans with PTSD, TBI, and comorbid TBI/PTSD. *Psychiatric Annals*, 39, 796–803.

Collins, K., & Bell, R. (1997). Personality and aggression: The dissipation-rumination scale. *Personality and Individual Differences*, 22, 751–755. doi:10.1016/S0191-8869(96)00248-6

Cooper, M. L., Agocha, V. B., & Sheldon, M. S. (2000). A motivational perspective on risky behaviors: The role of personality and affect regulatory processes. *Journal of Personality*, 68, 1059–1088. doi:10.1111/1467-6494.00126

Hayes, A. F. (2012). PROCESS: A versatile computational tool for observed variable moderation, mediation, and conditional process modeling. Retrieved from <http://www.afhayes.com/public/process2012.pdf>

Holman, E., & Silver, R. (1998). Getting “stuck” in the past: Temporal orientation and coping with trauma. *Journal of Personality and Social Psychology*, 74, 1146–1163. doi:10.1037/0022-3514.74.5.1146

Killgore W. D., Stetz M. C., Castro C. A., & Hoge C. W. (2008). The effects of prior combat experience on the expression of somatic and affective symptoms in deploying soldiers. *Journal of Psychosomatic Research*, 60, 379–385. doi:10.1016/j.jpsychores.2006.02.012

Kraaij, V., & Garnefski, N. (2006). The role of intrusion, avoidance, and cognitive coping strategies more than 50 years after war. *Anxiety, Stress & Coping: An International Journal*, 19, 1–14. doi:10.1080/10615800500412449

Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. W., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114, 163–173. doi:10.1016/j.jad.2008.06.026

Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspectives on Psychological Science*, 3, 400–424. doi:10.1111/j.1745-6924.2008.00088.x

Sezibera, V., Van Broeck, N., & Philippot, P. (2009). Intervening on persistent posttraumatic stress disorder: Rumination-focused cognitive and behavioral therapy in a population of young survivors of the 1994 genocide in Rwanda. *Journal of Cognitive Psychotherapy*, 23, 107–113. doi:10.1891/0889-8391.23.2.107

Shapiro, S. L., Oman, D., Thoresen, C. E., Plante, T. G., & Flinders, T. (2008). Cultivating mindfulness: Effects on well-being. *Journal of Clinical Psychology*, 64, 840–862. doi:10.1002/jclp.20491

Siegle, G. J., Moore, P. M., & Thase, M. E. (2004). Rumination: One construct, many features in healthy individuals, depressed individuals, and individuals with lupus. *Cognitive Therapy and Research*, 28, 645–668. doi:10.1023/B:COTR.0000045570.62733.9f

Thomsen, C. J., Stander, V. A., McWhorter, S. K., Rabenhorst, M. M., & Milner, J. S. (2011). Effects of combat deployment on risky and self-destructive behavior among active duty military personnel. *Journal of Psychiatric Research*, 45, 1321–1331. doi:10.1016/j.jpsychires.2011.04.003

Trapnell, P. D., & Campbell, J. D. (1999). Private self-consciousness and the five-factor model of personality: Distinguishing rumination from reflection. *Journal of Personality and Social Psychology*, 76, 284–304. doi:10.1037/0022-3514.76.2.284

Whitmer, A. J., & Banich, M. T. (2007). Inhibition versus switching deficits in different forms of rumination. *Psychological Science*, 18, 546–553. doi:10.1111/j.1467-9280.2007.01936.x